

University of Groningen

Development of multiscale simulation methods

Rzepliela, Andrzej Jerzy

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version

Publisher's PDF, also known as Version of record

Publication date:

2011

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):

Rzepliela, A. J. (2011). *Development of multiscale simulation methods*. s.n.

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

Development of multiscale simulation methods

Andrzej Rzepiela

- Cover:** Michał i Andrzej Rzepiela.
- Front:** Impression on a sampling of a configurational phase space in MD simulation by Michał.
- Back:** A coarse grained model and underlying ensemble of atomistic structures of a simple peptide molecule.

Printed by: Off Page, Amsterdam

© 2011 Andrzej Rzepiela

ISBN:978-94-6182-011-2

The research described in this thesis was carried out at the Groningen Biomolecular Sciences and Biotechnology Institute, Department of Biophysical Chemistry, University of Groningen, The Netherlands.



rijksuniversiteit
 groningen

Development of multiscale simulation methods

Proefschrift

ter verkrijging van het doctoraat in de
Wiskunde en Natuurwetenschappen
aan de Rijksuniversiteit Groningen
op gezag van de
Rector Magnificus, dr. E. Sterken,
in het openbaar te verdedigen op
maandag 12 september 2011
om 13.15 uur

door

Andrzej Jerzy Rzepiela

geboren op 1 oktober 1982
te Zielona Góra, Polen

Promotor: Prof. dr. S. J. Marrink

Beoordelingscommissie: Prof. dr. R. Broer
Prof. dr. P. G. Bolhuis
Prof. dr. N. van der Vegt

Contents

1	Introduction	1
1.1	Simulating the real world	2
1.2	The art of coarse-graining	3
1.2.1	Consistency conditions for CG models	3
1.2.2	Techniques to obtain CG models	5
1.2.3	Introducing dynamics into CG models	10
1.3	Multiscale algorithms	11
1.3.1	Multiscale atomistic – coarse-grained algorithms	12
1.4	Overview	14
2	Back mapping method	15
2.1	Abstract	16
2.2	Introduction	16
2.3	Methods	18
2.3.1	Initial placement of atomistic particles	18
2.3.2	Restrained simulated annealing	18
2.3.3	Release of the restraints	19
2.3.4	Reconstruction of solvent	19
2.4	Implementation	20
2.5	Simulation details	21
2.6	Example applications	22
2.6.1	NCQ peptide	22
2.6.2	DPPC and Cholesterol	28
2.6.3	Water	29
2.6.4	WALP peptide in solvated DPPC bilayer	31
2.7	Summary and Conclusions	33
2.8	Appendix: List of commands	35

3	Membrane poration by active peptides	37
3.1	Abstract	38
3.2	Introduction	38
3.3	Methods	40
3.3.1	Simulation protocol and force field	40
3.3.2	Resolution-exchange	42
3.3.3	System setup	42
3.4	Results and Discussion	42
3.4.1	Comparing the atomistic and coarse-grain pores	42
3.4.2	Characteristics of the CG pore	44
3.4.3	Effect of helicity on pore stability	46
3.4.4	Spontaneous pore formation in CG models	47
3.5	Summary and Conclusions	50
4	Fixed resolution hybrid method	53
4.1	Abstract	54
4.2	Introduction	54
4.3	Methods	56
4.3.1	Hybrid model	56
4.3.2	Coarse-grained potentials	56
4.3.3	Simulation setup	58
4.3.4	Analysis details	61
4.4	Results and Discussion	62
4.4.1	Comparing CG potentials for butane	62
4.4.2	PMFs for AA butane in CG solvent	63
4.4.3	Preferential solvation in mixed AA/CG butane	64
4.4.4	Association of AA dialanine in CG solvents	68
4.5	Summary and Conclusions	70
5	Hybrid “constant λ” method	73
5.1	Abstract	74
5.2	Introduction	74
5.3	Multiscaling theory	75
5.3.1	Derivation of forces	76
5.3.2	Temperature	78
5.4	Configurational entropy	80
5.4.1	Simulation details	80
5.4.2	Results for the MARTINI CG model	83
5.4.3	Results for the iterative Boltzmann model	86
5.5	Summary and Conclusions	87

<i>CONTENTS</i>	vii
6 Summary and Epilogue	89
Bibliography	95
Acknowledgments	111

List of Figures

1.1	Mapping of the MARTINI CG model.	4
1.2	Schematic representation of the AdResS multiscale method. . .	13
2.1	Ensemble of atomistic structures is mapped to a single CG representation: NCQ peptide.	22
2.2	Reconstruction of atomistic details of NCQ peptide.	25
2.3	Validation of the reconstruction procedure. Dihedral angle and potential energy distributions of NCQ peptide.	27
2.4	Reconstruction of a single DPPC molecule in vacuo.	29
2.5	Reconstruction of a single cholesterol molecule in vacuo. . . .	30
2.6	Reconstruction of a bulk SPC water.	31
2.7	Dipole moment distributions of restrained and free SPC water. .	32
2.8	Reconstruction of WALP20 in a solvated DPPC bilayer.	33
2.9	Helicity of WALP20 peptide as a function of annealing time. . .	34
3.1	Structure of the toroidal pore formed by magainin-H2 peptides in DPPC bilayers at multiple scales.	43
3.2	Density of water in the magainin pore.	45
3.3	Position of magainins in a toroidal pore.	46
3.4	Size of the magainin pore in function of the peptide helicity. . .	47
3.5	Radius of gyration of the magainin peptide in function of its helicity.	48
3.6	The time course of events when four magainin-H2 peptides are inserted into a DPPC bilayer modeled at the CG level.	49
3.7	The time course of events when fourteen magainin-H2 peptides are placed initially in water close to the surface of a short tail PC bilayer.	50
4.1	Graphical representation of the hybrid model.	57

4.2	Convergence of the iterative boltzmann procedure.	60
4.3	Coarse-grained butane–butane potentials.	63
4.4	Performance of different CG potentials in the hybrid model. . .	65
4.5	Hybrid model. Even mixture of AA and CG molecules.	66
4.6	Mixing of atomistic dialanine peptides in coarse-grained solvents.	69
4.7	Potential of mean force between two atomistic dialanines in CG solvent using MARTINI.	70
5.1	Effective potentials for CG hexadecane obtained with the itera- tive Boltzmann procedure.	81
5.2	Average configurational entropy of AA hexadecane in multiscale simulations for several values of λ . CG description of the system was represented with the MARTINI forcefield.	85
5.3	Average configurational entropy of AA hexadecane in multiscale simulations for several values of λ . CG description of the system was represented with the iterative Boltzmann procedure.	86

List of Tables

2.1	Parameters for reconstruction procedure.	21
4.1	Free energy of solvation of a CG butane molecule dissolved into its own bulk.	63
4.2	Preferential solvation parameters obtained from equal mixtures of AA and CG butane molecules.	67
5.1	Configurational entropies of hexadecane molecules, simulated with the multiscale protocol, according to Schlitter's quasi-harmonic approximation.	83
5.2	Observed scaling factors for Berendsen thermostats in simulations of liquid hexadecane at constant values of λ	84

